

Quantification of coral sperm collected during a synchronous spawning event

Aaron Teo¹, James R. Guest², Mei Lin Neo^{3,4}, Kareen Vicentuan³ and Peter A. Todd¹

¹ Department of Biological Sciences, National University of Singapore, Singapore

² SECORE International, Johor, Malaysia

³ Tropical Marine Science Institute, National University of Singapore, Singapore

⁴ St John's Island Marine Laboratory, National University of Singapore, Singapore

ABSTRACT

Most studies of coral reproductive biology to date have focused on oocyte numbers and sizes. Only one (*ex situ*) study has enumerated sperm numbers, even though these data have multiple potential applications. We quantified total coral sperm and eggs per gamete bundle collected from six species *in situ* during a synchronous spawning event in Singapore. Egg-sperm bundles were captured midwater as they floated towards the surface after being released by the colony. For each sample, a semi-transparent soft plastic bottle was squeezed and released to create a small suction force that was used to 'catch' the bundles. This technique provided several advantages over traditional methods, including low cost, ease of use, no diving prior to the night of collection needed, and the ability to target specific areas of the colony. The six species sampled were *Echinophyllia aspera*, *Favites abdita*, *F. chinensis*, *Merulina ampliata*, *M. scabricula* and *Platygyra pini*. The mean number of sperm packaged within one egg-sperm bundle ranged from 2.04×10^6 to 1.93×10^7 . The mean number of eggs per egg-sperm bundle ranged from 26.67 (SE \pm 3.27) to 85.33 (SE \pm 17.79). These data are critical for fertilisation success models, but the collection technique described could also be applied to studies requiring *in situ* spawning data at the polyp level.

Subjects Ecology, Marine Biology

Keywords Eggs, Coral, Gametes, Model, Sperm, Reproduction

INTRODUCTION

Approximately 86% of scleractinian coral species with documented reproductive traits have been identified as broadcast spawners (Baird, Guest & Willis, 2009). This mode of reproduction is prevalent in many aquatic taxa across multiple phyla (Leviton, 1998) and involves the shedding of gametes into the water column where external fertilisation and embryogenesis can occur (Szmant, 1986; Giese & Kanatani, 1987; Wray, 1995). The majority of broadcast spawning corals release simultaneously both eggs and sperm packaged together in buoyant egg-sperm bundles during a spawning event (Babcock & Heyward, 1986). These egg-sperm bundles aggregate on the surface, concentrating the gametes to ensure that fertilisation can occur in a small space relative to the entire three-dimensional seascape (Moláček, Denny & Bush, 2012).

Due to the multi-specific synchronous nature of many spawning events it is practically impossible to collect gametes from a particular species or colony from within spawn slicks.

Submitted 10 March 2016

Accepted 6 June 2016

Published 14 July 2016

Corresponding author

Peter A. Todd, dbspat@nus.edu.sg

Academic editor

James Reimer

Additional Information and
Declarations can be found on
page 6

DOI 10.7717/peerj.2180

© Copyright

2016 Teo et al.

Distributed under

Creative Commons CC-BY 4.0

OPEN ACCESS

Hence, studies that involve coral gamete collection have traditionally obtained spawned samples by trapping them before they reach the surface using nets placed directly over colonies ([Guest et al., 2010](#)) or by collecting them *ex situ* from colonies that have been moved to aquaria prior to spawning ([Negri & Heyward, 2001](#)). An alternative technique of collecting gamete data is via the histological analysis of polyp sections collected from excised coral fragments ([Nozawa & Lin, 2014](#)). As the development of eggs can be monitored within the ovaries, giving estimates of a polyp's fecundity, this approach has revealed abundant information regarding coral gametogenesis, spawning periods and synchronicity cues ([Manfred & Veghel, 1994](#); [Vermeij et al., 2004](#)), as well as mature oocyte sizes ([Nozawa & Lin, 2014](#)).

There is, however, a dearth of information on coral sperm compared to oocytes. Most studies focus on spermatogenesis using histological analyses ([Goffredo, Telò & Scanabissi, 2000](#)) to elucidate the impacts of anthropogenic disturbances such as sedimentation ([Gilmour, 1999](#); [Humphrey et al., 2008](#); [Jones, Ricardo & Negri, 2015](#)), heavy metal pollution ([Reichelt-Brushett & Harrison, 1999](#); [Negri & Heyward, 2001](#); [Reichelt-Brushett & Harrison, 2005](#); [Victor & Richmond, 2005](#)) and ocean acidification ([Morita et al., 2010](#)). Others have examined factors that influence fertilisation success ([Oliver & Babcock, 1992](#)), for example, changes in sperm motility due to increased acidity ([Morita et al., 2010](#)) as well as damage and clumping due to suspended sediment ([Ricardo et al., 2015](#)). Sperm data, such as the amount released during a spawning, are mandatory in simulation models that attempt to make realistic predictions of sperm dilution and fertilisation rates ([Babcock, Mundy & Whitehead, 1994](#); [Metaxas, Scheibling & Young, 2002](#)). Such models allow for general or site-specific estimates of a population's reproductive output—complementing other forms of data used in reef health monitoring and assessment, e.g. life tables and recruitment rates, thereby contributing to conservation management plans.

Most notably, other than for *Acropora digitifera* ([Paxton et al., 2015](#)), there is a conspicuous absence of data regarding the number of sperm released by scleractinian corals during spawning. Hence, the main aim of the present study was to quantify the amount of sperm within a coral gamete bundle through *in situ* sampling of various coral species during a synchronous spawning event using a novel yet simple technique of collecting small samples of intact coral gamete bundles in midwater.

MATERIALS AND METHODS

Study site

Pulau Satumu (Raffles Lighthouse) is a small rocky island measuring only 1.3 ha. Due to the relatively strong currents and distance (~14 km) from Singapore's main island, the reefs around Pulau Satumu are less affected by chronic sedimentation compared to other local reefs. Multi-species coral spawning events occur during the week following the March or April full moon ([Guest et al., 2005](#)).

Gamete collection

Coral gamete bundles were collected during the spawning events that occurred on the nights of 19th and 20th April 2014 between 20.00–22.00 h from the fringing reefs to the west of Pulau Satumu. Opportunistic sampling within a 50 × 5 m² belt transect

was conducted by identifying colonies that were ready to spawn (by observing gamete bundles “setting” just below the polyp mouth) and waiting for the release of the gametes.

Using an underwater digital camera, each selected colony was photographed with a uniquely labeled transparent ziplock bag placed alongside it so that the bags could later be matched to the coral species. Semi-transparent 30 ml soft plastic bottles (outer dimensions: 59 × 27 mm, length × diameter; 10 mm diameter nozzle opening) were used to collect gamete bundles. The bottle was squeezed and released to create a small suction force that was used to ‘catch’ bundles as they floated towards the surface after being released by the colony. This technique enabled the capture of intact gamete bundles from specific colonies. The bottles were handled carefully to avoid breaking the bundles prematurely. On the first day three bundles were captured within each bottle from five species whereas on the second day six bundles were captured within one bottle from one species.

Three bottles of samples were collected per colony and placed in their colony-specific ziplock bags for transportation back to the boat. The intact gamete bundles and seawater contained within each bottle were transferred into labeled falcon tubes prefilled with sufficient formaldehyde to produce 50 ml of solution at 3.7% concentration. The falcon tubes were then agitated by hand to break apart the gamete bundles and fix the eggs and sperm released.

Gamete quantification

After the eggs were filtered out using an 80 µm sieve, three 10 µl aliquots were extracted per (agitated) falcon tube with a micropipette. Each aliquot was stained with 10 µl of trypan blue and left to stand for 15 min before the contents of the aliquot was injected into a Neubauer haemocytometer for counting. Sperm were visually distinguished from other matter in the solution by the distinctive shape of the sperm heads. All sperm within the 25 large squares (0.1 mm³ total volume) in the center grid were counted. The filtered eggs for each falcon tube were photographed and then counted using ImageJ (ImageJ 1.48 v; National Institutes of Health, Bethesda, MD, USA).

RESULTS AND DISCUSSION

Six species (*Echinophyllia aspera*, *Favites abdita*, *F. chinensis*, *Merulina ampliata*, *M. scabricula*, and *Platygyra pini*) were sampled successfully. The mean number of sperm packaged within one egg-sperm bundle ranged from 2.04×10^6 to 1.93×10^7 . The greatest number of sperm was released by *P. pini*, whereas the least was from *F. abdita* (Table 1). The mean (± SE) number of eggs per egg-sperm bundle varied from 26.67 (SE ± 3.27) to 85.33 (SE ± 17.79). *F. chinensis* had the highest sperm: egg ratio whereas *F. abdita* had the lowest (Table 1). There was generally a positive relationship between the number of eggs and number of sperm per bundle (Fig. 1).

The technique of using small transparent soft plastic bottles as collection devices was effective at capturing individual gamete bundles and was simple enough to be used one-handed (freeing up the other hand to hold a torch, for example). Unlike the

Table 1 Mean (\pm SE) number of eggs and sperm per bundle for the seven colonies sampled in April 2014. Three samples (bottles) of egg-sperm bundles were collected per colony. Two colonies were sampled for *Echinophyllia aspera* (labelled 1 and 2); for all other species only one colony was sampled.

Species	Date	No of bundles in sample	Mean eggs (bundle ⁻¹)	Mean sperm ($\times 10^6$ bundle ⁻¹)	Sperm-egg ratio
<i>Favites abdita</i>	19/04	3	48.33 \pm 5.01	2.04 \pm 0.30	4.2 ($\times 10^4$):1
<i>F. chinensis</i>	19/04	3	26.67 \pm 3.27	9.33 \pm 1.62	35.0 ($\times 10^4$):1
<i>Merulina ampliata</i>	19/04	3	72.78 \pm 9.03	8.44 \pm 2.03	11.6 ($\times 10^4$):1
<i>M. scabricula</i>	19/04	3	70.67 \pm 7.65	4.33 \pm 0.72	6.1 ($\times 10^4$):1
<i>Platygyra pini</i>	19/04	3	85.33 \pm 17.79	19.33 \pm 2.85	22.7 ($\times 10^4$):1
<i>Echinophyllia aspera</i> (colony 1)	20/04	6	41.67 \pm 25.36	5.76 \pm 2.37	13.8 ($\times 10^4$):1
<i>E. aspera</i> (colony 2)	20/04	6	60.27 \pm 21.72	12.63 \pm 1.31	21.0 ($\times 10^4$):1

traditional approaches of using traps positioned above selected specimens or collecting entire colonies and moving them to tanks on land or on a boat, our technique required little preparation and no prior diving to set up traps or collect colonies. No aquarium facilities were required and no damage was done to donor colonies. Critically, the technique enables the direct quantification of sperm released during actual spawning. While this information may be extracted from egg-sperm bundles captured in traps positioned over colonies, or with nets swept by hand, there remains the possibility that these become broken up. Variation in polyp fecundity due to polyp position (Nozawa & Lin, 2014) is also not distinguishable when using trapped samples (as they are mixed during collection).

We required sperm quantity data for an ongoing modelling project to predict the fertilisation success of synchronous spawning corals and such data are likely to be important for other contemporary and future fertilisation models. However, additional information could be extracted from sperm collected using this technique. For example, sperm number and stage of development is usually described by the size of the spermaries present in tissue samples (Nozawa & Lin, 2014) without any quantification of the number of competent sperm present. The scarcity of sperm count data can hinder other aspects of coral reproduction research, including assessing the risks of sperm limitation and its antithesis, polyspermy, as well as the reproductive strategies that may have evolved to cope with these potential issues. Morphological data for sperm are also scarce, with few studies detailing their characteristics (Harrison, 1985; Steiner, 1991; Hagedorn et al. 2006) despite the possibility that these can yield information on sexual reproduction patterns (Harrison, 1985).

There was substantial variation in the average number of gametes per bundle sampled, with differences of up to ten times more sperm between samples recorded (after accounting for the number of bundles per sample). These differences were found across all levels, ranging from species to individual colonies and even individual samples of bundles from the same colony. The observed differences are likely to be due to a combination of factors such as species and genotypic differences in reproductive output, as well as the influence of age (Sakai, 1998), colony and polyp size (Hall & Hughes, 1996;




Figure 1 Number of sperm per bundle for the seven colonies sampled in April 2014. Each point represents the mean values for three egg-sperm bundles collected in one bottle (six egg-sperm bundles per bottle for *Echinophyllia aspera*). Three bottles were collected per colony. Two colonies were sampled for *E. aspera* (labelled 1 and 2); for all other species only one colony was sampled.

Sakai, 1998; Nozawa & Lin, 2014) and polyp position within the colony (Sakai, 1998; Nozawa & Lin, 2014). Future studies could use the technique described here to target specific polyps in a colony to better understand intra-colonial variation in gamete output. Although this can be achieved via histology (and the breaking off of fragments from around the colony) not all gametes may be released during spawning as they can reabsorbed by the polyp (Harrison & Wallace, 1990), providing erroneous data. As the technique is non-destructive, repeatedly examining the same polyp also becomes a possibility.

As the threat of extinction looms over a third of all reef-building coral species (Carpenter et al., 2008; Plaisance et al., 2011), the need to improve the efficiency and effectiveness of management and restoration plans is increasingly urgent. Reproductive output data are critical to such conservation efforts; however, existing knowledge gaps hinder coral fertilisation models and the ability to predict the impact of disturbances. Our study introduces a simple, effective and non-destructive means of sampling intact egg-sperm bundles during coral spawning in the field. The technique helped elucidate the number of sperm packaged within an egg-sperm bundle and can potentially capture polyp-level variation in fecundity.

ACKNOWLEDGEMENTS

Many thanks to Patrick Cabaitan and Danwei Huang for help with coral identification as well as Lutfi Afiq Rosli and Hai Xin Loke for help with collecting the gamete bundles.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

The National Parks Board, Singapore, funded the research vessel used during the spawning event. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests

The authors declare that they have no competing interests.

Author Contributions

- Aaron Teo conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.
- James R. Guest wrote the paper, reviewed drafts of the paper.
- Mei Lin Neo performed the experiments, reviewed drafts of the paper.
- Kareen Vicentuan performed the experiments, reviewed drafts of the paper.
- Peter A. Todd conceived and designed the experiments, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.

Data Deposition

The following information was supplied regarding data availability:

The raw data has been supplied as [Supplemental Dataset Files](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.2180#supplemental-information>.

REFERENCES

- Babcock RC, Heyward AJ. 1986. Larval development of certain gamete-spawning scleractinian corals. *Coral Reefs* 5(3):111–116 DOI 10.1007/BF00298178.
- Babcock RC, Mundy CN, Whitehead D. 1994. Sperm diffusion models and in situ confirmation of long-distance fertilization in the free-spawning asteroid *Acanthaster planci*. *Biological Bulletin* 186(1):17–28 DOI 10.2307/1542033.
- Baird AH, Guest JR, Willis BL. 2009. Systematic and biogeographical patterns in the reproductive biology of scleractinian corals. *Annual Reviews of Ecology, Evolution and Systematics* 40(1):551–571 DOI 10.1146/annurev.ecolsys.110308.120220.
- Carpenter KE, Abrar M, Aeby G, Aronson RB, Banks S, Bruckner A, Chiriboga A, Cortés J, Delbeek JC, DeVantier L, Edgar GJ, Edwards AJ, Fenner D, Guzmán HM, Hoeksema BW, Hodgson G, Johan O, Licuanan WY, Livingstone SR, Lovell ER, Moore JA, Obura DO, Ochavillo D, Polidoro BA, Precht WF, Quibilan MC, Reboton C, Richards ZT, Rogers AD, Sanciangco J, Sheppard A, Sheppard C, Smith J, Stuart S, Turak E, Veron JEN, Wallace C, Weil E, Wood E. 2008. One-third of reef-building corals face elevated extinction risk from climate change and local impacts. *Science* 321(5888):560–563 DOI 10.1126/science.1159196.

- Giese AC, Kanatani H. 1987. Maturation and spawning. In: Giese AC, Pearse JS, Pearse VB, eds. *Reproduction of Marine Invertebrates Vol. IX: Seeking Unity in Diversity*. Palo Alto: Blackwell Scientific, 251–329.
- Gilmour J. 1999. Experimental investigation into the effects of suspended sediment on fertilisation, larval survival and settlement in a scleractinian coral. *Marine Biology* 135(3):451–462 DOI 10.1007/s002270050645.
- Goffredo S, Telò T, Scanabissi F. 2000. Ultrastructural observations of the spermatogenesis of the hermaphroditic solitary coral *Balanophyllia europaea* (Anthozoa, Scleractinia). *Zoomorphology* 119(4):231–240 DOI 10.1007/PL00008495.
- Guest JR, Baird AH, Goh BPL, Chou LM. 2005. Reproductive seasonality in an equatorial assemblage of scleractinian corals. *Coral Reefs* 24(1):112–116 DOI 10.1007/s00338-004-0433-7.
- Guest JR, Heyward A, Omori M, Iwao K, Morse A, Boch C. 2010. Rearing coral larvae for reef rehabilitation. In: Edwards AJ, ed. *Reef Rehabilitation Manual*. St Lucia: The Coral Reef Targeted Research & Capacity Building for Management Program, 73–98. Available at http://www.gefcoral.org/Portals/53/downloads/Reef%20Rehabilitation%20Manual_web.pdf (accessed 1 March 2016).
- Hagedorn M, Carter VL, Steyn RA, Krupp D, Leong JC, Lang RP, Tiersch TR. 2006. Preliminary studies of sperm cryopreservation in the mushroom coral, *Fungia scutaria*. *Cryobiology* 52(3):454–458 DOI 10.1016/j.cryobiol.2006.03.001.
- Hall VR, Hughes TP. 1996. Reproductive strategies of modular organisms: comparative studies of reef-building corals. *Ecology* 77(3):950–963 DOI 10.2307/2265514.
- Harrison PL. 1985. Sexual characteristics of scleractinian corals: systematic and evolutionary implications. *Proceedings of the Fifth International Coral Reef Congress, 27 May–1 June 1985*, Vol. 4, Tahiti: Symposia and Seminars (B), 337–342.
- Harrison PL, Wallace CC. 1990. Reproduction, dispersal and recruitment of scleractinian corals. In: Dubinsky Z, ed. *Ecosystems of the World 25: Coral Reefs*. Amsterdam: Elsevier Science, 133–207.
- Humphrey C, Weber M, Lott C, Cooper T, Fabricius K. 2008. Effects of suspended sediments, dissolved inorganic nutrients and salinity on fertilisation and embryo development in the coral *Acropora millepora* (Ehrenberg, 1834). *Coral Reefs* 27(4):837–850 DOI 10.1007/s00338-008-0408-1.
- Jones R, Ricardo GF, Negri AP. 2015. Effects of sediments on the reproductive cycle of corals. *Marine Pollution Bulletin* 100(1):13–33 DOI 10.1016/j.marpolbul.2015.08.021.
- Levitan DR. 1998. 6 Sperm limitation, gamete competition and sexual selection in external fertilizers. In: Birkhead TR, Møller AP, eds. *Sperm Competition and Sexual Selection*. San Diego: Academic Press, 175–218.
- Manfred LJ, Veghel V. 1994. Reproductive characteristics of the polymorphic Caribbean reef building coral *Montastrea annularis*. Gametogenesis and spawning behavior. *Marine Ecology Progress Series* 109:209–219 DOI 10.3354/meps109209.
- Metaxas A, Scheibling RE, Young CM. 2002. Estimating fertilization success in marine benthic invertebrates: a case study with the tropical sea star *Oreaster reticulatus*. *Marine Ecology Progress Series* 226:87–101 DOI 10.3354/meps226087.
- Moláček J, Denny M, Bush JWM. 2012. The fine art of surfacing: its efficacy in broadcast spawning. *Journal of Theoretical Biology* 294:40–47 DOI 10.1016/j.jtbi.2011.10.013.
- Morita M, Suwa R, Iguchi A, Nakamura M, Shimada K, Sakai K, Suzuki A. 2010. Ocean acidification reduces sperm flagellar motility in broadcast spawning reef invertebrates. *Zygote* 18(2):103–107 DOI 10.1017/S0967199409990177.

- Negri AP, Heyward AJ. 2001. Inhibition of coral fertilisation and larval metamorphosis by tributyltin and copper. *Marine Environmental Research* 51(1):17–27 DOI 10.1016/S0141-1136(00)00029-5.
- Nozawa Y, Lin C-H. 2014. Effects of colony size and polyp position on polyp fecundity in the scleractinian coral genus *Acropora*. *Coral Reefs* 33(4):1057–1066 DOI 10.1007/s00338-014-1185-7.
- Oliver J, Babcock R. 1992. Aspects of the fertilization ecology of broadcast spawning corals: sperm dilution effects and in situ measurements of fertilization. *Biological Bulletin* 183(3):409–417 DOI 10.2307/1542017.
- Paxton CW, Baria MVB, Weis VM, Harii S. 2015. Effect of elevated temperature on fecundity and reproductive timing in the coral *Acropora digitifera*. *Zygote* 24(4):511–516 DOI 10.1017/S0967199415000477.
- Plaisance L, Caley MJ, Brainard RE, Knowlton N. 2011. The diversity of coral reefs: what are we missing? *PLoS ONE* 6(10):e25026 DOI 10.1371/journal.pone.0025026.
- Reichelt-Brushett AJ, Harrison PL. 1999. The effect of copper, zinc and cadmium on fertilization success of gametes from scleractinian reef corals. *Marine Pollution Bulletin* 38(3):182–187 DOI 10.1016/S0025-326X(98)00183-0.
- Reichelt-Brushett AJ, Harrison PL. 2005. The effect of selected trace metals on the fertilization success of several scleractinian coral species. *Coral Reefs* 24(4):524–534 DOI 10.1007/s00338-005-0013-5.
- Ricardo GF, Jones RJ, Clode PL, Humanes A, Negri AP. 2015. Suspended sediments limit coral sperm availability. *Scientific Reports* 5:18084 DOI 10.1038/srep18084.
- Sakai K. 1998. Effect of colony size, polyp size, and budding mode on egg production in a colonial coral. *Biological Bulletin* 195(3):319–325 DOI 10.2307/1543143.
- Steiner SCC. 1991. Sperm morphology of scleractinians from the Caribbean. *Hydrobiologia* 216(1):131–135 DOI 10.1007/BF00026453.
- Szmant AM. 1986. Reproductive ecology of Caribbean reef corals. *Coral Reefs* 5(1):43–54 DOI 10.1007/BF00302170.
- Vermeij MJA, Sampayo E, Bröker K, Bak RPM. 2004. The reproductive biology of closely related coral species: gametogenesis in *Madracis* from the southern Caribbean. *Coral Reefs* 23(2):206–214 DOI 10.1007/s00338-004-0368-z.
- Victor S, Richmond RH. 2005. Effect of copper on fertilization success in the reef coral *Acropora surculosa*. *Marine Pollution Bulletin* 50(11):1448–1451 DOI 10.1016/j.marpolbul.2005.09.004.
- Wray GA. 1995. Evolution of larvae and developmental modes. In: McEdward LR, ed. *Ecology of Marine Invertebrate Larvae*. Florida: CRC Press, 413–448.